



## ACUTE NUMBER ONE REPAIR OF EXTRAARTICULAR LIGAMENTS AND STAGED SURGERY IN MORE THAN ONE LIGAMENT KNEE INJURIES

**Mohammad  
Imdadul Hoque  
Shakil**

**Nahida Islam**

Orthopedic Consultant, Department of Orthopedic, UHC sadar, Mymensingh, Bangladesh

Associate Professor, Head Of Department, Department of Dermatology and Venereology, Community Based medical College & Hospital, Bangladesh.

### ABSTRACT

**Background:** The motive of this observe is to compare the consequences of acute number one repair of extraarticular ligaments with staged surgical treatment for acute knee dislocations (KDs) and multiligament knee injuries (MLKIs). **Materials and methods** The study has conducted in Department of Orthopedic, UHC sadar, Mymensingh. During the period of January 2013 and May 2021, 67 consecutive patients identified with MLKI or KD have been noted or visited our institution. Of these, 37 sufferers who underwent acute restore of extraarticular ligaments inside three weeks of damage had been included on this observe. These patients were retrospectively categorised into two corporations: people who underwent best primary repair (repair group) and people who underwent staged reconstructive surgical operation (staged group). Follow-up exam protected range of motion (ROM), knee joint balance (Lachman test, posterior drawer take a look at, and varus and valgus stress test), Lysholm knee score, Tegner interest scale, and Knee damage and Osteoarthritis and effect Score (KOOS). **Results:** Twelve of the 37 sufferers did not want or preference similarly surgical treatment and were protected inside the restore institution. No great distinction become determined in demographic facts among the restore and staged businesses. Although staged surgery decreased high-quality posterior drawer check results, no giant difference became located between the 2 businesses concerning ROM, different knee joint balance tests, Lysholm rankings, Tegner scale, or KOOS. **Conclusions:** all patients lower back to their activities of each day residing and preinjury career ranges. Acute primary restore of extraarticular ligaments gives critical knee stability with out varus/valgus instability and may lessen the need for next cruciate ligament reconstruction.

### KEYWORDS

Knee dislocations, Multiligament knee injuries, Primary repair, Staged surgery

#### Introduction:

Knee dislocations (KDs) and multiligament knee injuries (MLKIs) are extreme knee traumas which involve intra- and extraarticular ligament tears, often with concomitant vascular and nerve damage and a fracture across the knee. Because popliteal artery lesion is a limb-threatening injury, early revascularization have to be prioritized to avoid limb amputation [1, 2]. Concomitant different organ traumas, along with open fracture and head trauma, may also in addition compromise the foremost timing of MLKI treatment. Therefore, it's far hard to apply a unmarried technique and best surgical timing. Since MLKIs and KDs are uncommon and often heterogeneous, as cited above, minimal proof is to be had, resulting in a loss of consensus concerning the best remedy [3, 4]. Although conservative and surgical treatments had been stated, surgical interventions have typically been endorsed due to bad outcomes after conservative remedy [5,6,7]. Currently, conservative remedy is completely selected for patients who are unfit for surgical treatment, frail, or sedentary [7].

Surgical intervention varies from the primary repair of damaged ligaments to anatomical ligament reconstruction in both a simultaneous or staged fashion [5, 8]. Early surgical treatment has been endorsed to improve outcomes [9,10,11], and the crucial time to reestablish anatomic relationships is the first three weeks after harm [12]. Acute ligament reconstruction improves postoperative knee stability [13] and might boom the rate of arthrofibrosis, which causes deterioration in knee function and calls for additional surgical procedures [14]. On the opposite, delayed reconstruction may additionally offer the time for natural healing of extraarticular ligaments and reduce postoperative arthrofibrosis [5, 15, 16]; it requires multiple grafts and tunnels for reconstructions, resulting in donor-site morbidity and menace of excavate union [17]. Staged surgery, which entails restore of the extraarticular ligaments in the intense level and subsequent reconstruction of the cruciate ligaments at a later degree, showed incredible scientific effects [18]. However, staged surgical operation calls for a couple of surgical procedures, and

this prolongs rehabilitation. Recently, the number one repair of knee ligaments, together with the intraarticular ligaments, has attracted interest as it has the benefit of maintaining the local tissues and fending off the want for graft harvesting or greater invasive surgical procedure [19,20,21].

In our experience of MLKI treatments, a few patients did no longer undergo right primary restore due to polytrauma, requiring prolonged extensive care or revascularization surgery for popliteal arterial damage, and had large residual knee instability in spite of undergoing not on time reconstruction. After experiencing those instances, we changed our surgical approach to early restore of the extraarticular ligaments, specifically the posterior pill structure, on the time of revascularization surgical operation. Intraarticular cruciate ligament reconstructions are done if the affected person goals further surgical operation. The motive of this look at is to compare the consequences of acute number one repair of extraarticular ligaments with staged surgical operation in acute KDs and MLKIs. We hypothesize that ideal primary repair of extraarticular ligaments not only improves the consequences of intraarticular cruciate ligament reconstruction but additionally reduces the frequency of cruciate ligament reconstruction.


#### Patients and methods

Patients: In this take a look at, MLKIs have been defined as disruption of as a minimum two of the four important knee ligament structures [anterior cruciate ligament (ACL), posterior cruciate ligament (PCL), posteromedial corner (PMC), and posterolateral corner (PLC)] [5]. The study has conducted in Department of Orthopedic, UHC sadar, Mymensingh. During the period of January 2013 and May 2021, a consecutive series of 67 patients recognized with MLKI or KD have been cited or visited our organization. Inclusion standards had been (1) radiographically documented KD; (2) PCL harm with associated accidents to the PMC, inclusive of the medial collateral ligament (MCL), and/or associated injuries to PLC, including the lateral

### \*CORRESPONDING AUTHOR Nahida Islam\*

Associate Professor, Head Of Department, Department of Dermatology and Venereology, Community Based medical College & Hospital, Bangladesh

collateral ligament (LCL); (3) bicruciate ligament injury and associated harm to as a minimum one collateral ligament (KD-IIIM or KD-IIIL) [22]; and (4) injury to all four major ligaments (KD-IV). On the contrary, exclusion standards had been (1) persistent MLKIs or KDs, [2] no acute number one restore due to extended intensive care, [3] open knee dislocation, [4] knees with osteoarthritis, [5] patients with ACL harm and grade III MCL injury who underwent simultaneous ACL reconstruction and MCL repair, and [6] failure to finish the study questionnaire (Fig. 1). 37 sufferers who underwent acute number one repair of extraarticular ligaments inside 3 weeks after injury met the inclusion standards and shaped the study organization (Table 1). The common age at damage become 48.6 ± 21.3 years (14–70 years), and there had been 26 men and 11 girls. The mechanisms of injury were 18 high-electricity traumas, such as site visitors injuries and falls from heights greater than 2 m, 12 sports-related accidents, and 7 other low-strength traumas. These sufferers had been categorised into two organizations: people who underwent most effective primary restore (repair group) and those who underwent staged surgery (staged organization). After approval from our institution's ethics committee, all patients supplied knowledgeable written consent earlier than inclusion within the look at.



Flowchart of study. KDs: knee dislocations; MLKIs: multiligament knee injuries


**Table 1 Patient demographics From: Acute primary repair of extraarticular ligaments and staged surgery in multiple ligament knee injuries**

	All injured knees (N = 37)	Repair group (N = 11)	Staged group (N = 26)	p-Value
Age (years): mean ± SD (range)	48.0 ± 20.6 (14–75)	55.9 ± 17.5 (18–75)	42.9 ± 21.2 (14–73)	0.093
Sex (male : female)	26 (67.7%):10 (32.3%)	9 (75.0%):3 (25.0%)	12 (63.2%):7 (36.8%)	0.697
BMI (kg/m <sup>2</sup> ): mean ± SD (range)	25.1 ± 4.4 (19.7–39.5)	25.7 ± 5.3 (20.8–39.5)	24.8 ± 3.8 (19.7–32.0)	0.703
Time to primary repair (days)	5.9 ± 5.5 (0–20)	6.3 ± 7.1 (0–20)	5.7 ± 4.5 (0–14)	0.646
Damaged ligaments N (%)				
PCL, PMC, and/or PLC	6 (19.3%)	3 (25.0%)	3 (15.8%)	0.879
ACL, PCL, MCL (KD III-M)	18 (58.1%)	7 (58.3%)	11 (57.9%)	
ACL, PCL, LCL (KD III-L)	3 (9.7%)	1 (8.3%)	2 (10.5%)	
ACL, PCL, MCL, LCL (KD IV)	4 (12.9%)	1 (8.3%)	3 (15.8%)	
Associated injuries, N (%)				
Nerve injury	3 (9.7%)	1 (8.3%)	2 (10.5%)	1.000
Vascular injury	5 (16.1%)	3 (25.0%)	2 (10.5%)	0.350

Follow-up (months) mean ± SD (range)	60.9 ± 31.7 (24–160)	50.0 ± 23.0 (24–78)	67.6 ± 34.9 (24–160)	0.164
--------------------------------------	----------------------	---------------------	----------------------	-------

1. SD standard deviation, N number, BMI body mass index, ACL anterior cruciate ligament, PCL posterior cruciate ligament, PMC posteromedial corner, PLC posterolateral corner, MCL medial collateral ligament, LCL lateral collateral ligament, KD knee dislocation.

**Preoperative exam:** After management, radiographic critiques, inclusive of magnetic resonance imaging (MRI), had been performed as quickly as viable to decide surgical techniques (Fig. 2). Computed tomography (CT) angiography turned into usually carried out if the patient showed any suspected signs of popliteal arterial injury, together with ankle-brachial strain index > zero.Nine. After examination of knee instability beneath anesthesia, diagnostic arthroscopy became quick carried out to assess associated intraarticular lesions in all sufferers besides for those with vascular damage. Meniscal lesions, inclusive of locked meniscus, were handled arthroscopically if discovered.



**Fig. 2 From: sensitive major repair of extraarticular ligaments and thesian surgery in multiple ligament knee injuries**


MRI and CT angiography and assessment beneath anesthesia. An 18-12 months-antique baseball player who suffered multiligament knee accidents in the course of a three-hitter. A Sagittal MRI showing PCL damage and posterior capsular injury; b coronal MRI revealing avulsion of the posterolateral ligament complicated from the fibular head; c angio-CT revealing occlusion of the popliteal artery; d, e evaluation beneath anesthesia displaying intense posterior and varus instability.

**Acute number one restore:**

Patients were positioned in supine position, and extraarticular medial-sided and lateral-sided accidents had been repaired. Through medial or lateral longitudinal incision, broken systems were cautiously identified. The injured collateral ligament become sutured using pull-out sutures (no. 2 Ethibond; Ethicon, Somerville, NJ) from the intact attachment closer to the avulsed ends. Subsequently, the avulsed ends have been constant to their anatomical insertion the usage of smooth suture anchors underpulling the Ethibond sutures. Similarly, the pull-out sutures have been used to reinforce the fixation of the MCL or LCL to the encompassing smooth tissues. The systems of the PMC and PLC have been additionally anatomically constant to their anatomical web site of insertion the use of gentle suture anchors. All accidents of the tablet across the joint have been treated with number one repair using absorbable sutures (2-0 Vicryl; Ethicon) and smaller suture anchors to offer fixation factors. If there have been any avulsion fractures continuous with the ligaments, the fragment was constant by using screws. Since posterior systems of the knee grow to be taut in an extension position, restore of those structures was completed with the knee held in extension. After number one restore, we confirmed whether the knee can be completely extended.

If the patient suffered popliteal arterial harm, emergent vascular surgical procedure turned into usually done to save you limb amputation. At our institution, orthopedic hand surgeons finished those vascular reconstructions, along with primary arterial sutures and a reverse saphenous vein graft. Patients had been positioned in prone position, and their knees were barely flexed (Figs. 3, 4). Through the posterior crank skin incision, vascular surgical treatment was achieved; and in the end, acute number one upkeep of extraarticular


ligaments have been achieved thru the identical incision. For patients with peroneal nerve palsy, nerve release turned into completed, followed by way of number one repairs of extraarticular ligaments.



**Fig. 3 From: sensitive major repair of extraarticular ligaments and thespian surgery in multiple ligament knee injuries**

Vascular surgical procedure and number one repair of posterolateral complicated of left knee (identical patient as in Fig. 2). A Skin incision for posterior approach; b popliteal artery thrombosed due to intimal rupture (arrows); c reversed saphenous vein graft (arrows), d PLC (arrow heads) avulsed from fibular head (arrow); e suture anchors inserted into fibular head; and f PLC constant by using suture anchors and torn posterior tablet repaired.

Postoperative radiograph (same affected person as in Fig. 2). A, b Radiograph taken 4 years after number one restore showing no full-size alternate in osteoarthritis; c, d posterior sag view of bilateral knee. Radiograph of right knee suggests posterior laxity of thirteen mm. This patient does now not need similarly surgical procedure and has again to recreational baseball play without restriction.



**From:4 sensitive major restore of extraarticular ligaments and thespian surgical treatment in more than one ligament knee injuries**


#### Postoperative rehabilitation after primary restore:

Postoperatively, the patient's knee modified into fixed externally with a brace. Patients commenced isometric muscle-strengthening sports activities, consisting of patella setting and right away leg elevating, the day after surgical treatment. If viable, sufferers were allowed non-weight-bearing gait with crutches as quickly as feasible. Range of movement (ROM) carrying activities the use of non-stop passive motion devices and partial weight-bearing gait became commenced at the start of the 0.33 postoperative week. Patients progressed to full ROM workout and full weight-bearing gait after 6 weeks. No open chain physical video games were allowed for the primary three months.

Staged reconstructionACruciate ligament reconstruction turn out to be typically encouraged for more youthful, active sufferers as an elective surgical operation after number one restore. Once the affected character had gained a sufficient ROM in their knees, they underwent staged surgery, typically approximately 6 months after number one restore. Cruciate ligament reconstruction turned into performed the use of a double-package deal method with an ipsilateral autogenous hamstring tendon (Fig. Five). When simultaneous double-package ACL and PCL reconstructions were done, contralateral hamstring tendons have been moreover harvested. If the affected person did now not preference in addition surgical procedure, the surgical remedy modified into completed entirely with acute primary repair. After staged surgical treatment, practically similar rehabilitation as after the number one repair changed into done.

Double-package ACL and PCL reconstructions of right knee (26-yr-vintage male judoist). A preoperative MRI displaying torn ACL and PCL; b, c postoperative MRI displaying reconstructed ACL (arrow

heads) and PCL (arrows), d arthroscopic view of double-package ACL and PCL reconstructions, and e, f postoperative radiograph. ACL: anterior cruciate ligament; PCL: posterior cruciate ligament; MRI: magnetic resonance imaging



**Fig. 5**

**Outcome checks:** Postoperative headaches, which include contamination, had been assessed. Postoperative ROM and knee stability were assessed at final take a look at-up. Knee stability at very last examine-up turned into defined with the useful resource of a Lachman test  $\leq$  grade I, a posterior drawer take a look at  $\leq$  grade II, and varus and valgus instability  $\leq$  grade II. The Lysholm knee rating modified into used for assessment of every day purposeful hobby at final observe-up, and activity stages at preinjury and final observe-up were evaluated with the Tegner pastime scale. Subjective pride became assessed using the Knee Injury and Osteoarthritis and Outcome Score (KOOS) [23]. These results have been as compared a number of the restore and staged businesses.

**Statistical analysis:** Descriptive records are supplied as suggest  $\pm$  standard deviation for non-stop variables, and the Tegner interest scale became calculated with median and variety values. Preoperative demographics, postoperative ROM, and medical rankings, which includes the Tegner hobby scale, the Lysholm score, and the KOOS, have been as compared a number of the repair and staged organizations the use of the Mann-Whitney U take a look at. In addition, kind of knee dislocation, price of concomitant peroneal nerve or vascular damage, and presence of knee instability had been as compared many of the 2 groups using a chi-rectangular take a look at or Fisher's real test. Data enter and analyses had been completed the use of SPSS model 25.0 I(SPSS Inc., Chicago, IL). P-Value  $<$  0.05 grow to be considered statistically big.

**Results:** Twelve of the 37 sufferers did now not need or want similarly surgical procedure and had been protected inside the restore organization (38.7%). The other 19 patients underwent cruciate ligament reconstruction and have been covered inside the staged organization (61.3%). The common age inside the restore group turned into fifty five.9  $\pm$  17.5 (18-seventy five) years, and there were 9 men and 3 ladies. The common age in the staged organisation have become 42.9  $\pm$  21.2 (14-73) years, and there had been 12 men and seven women. The follow-up durations inside the restore and staged agencies have been 50.0  $\pm$  23.0 (24-78) months and 67.6  $\pm$  34.9 (24-a hundred and sixty) months, respectively. No large difference turned into positioned in patient traits among the restore and staged corporations (Table 1). The maximum not unusual harm changed into a combination of ACL, PCL, and MCL (KD III-M) occurring with a frequency of 58.1%.

No big headaches had been positioned after both number one restore or staged surgery. One affected character had pores and skin necrosis requiring unfastened pores and pores and skin grafting after popliteal arterial surgical remedy in the number one restore group. Regarding medical effects at final study-up, no huge difference end up placed among the two groups regarding ROM or knee balance, besides posterior instability ( $p = 0.006$ ) (Table 2). No sufferers offered postoperative grade III varus or valgus instability in both enterprise. Postoperative Tegner hobby scales reduced compared with preinjury ones in every businesses, and all patients returned to their activities of daily living and preinjury occupational diploma. Lysholm score, Tegner interest scale, and all subscales of the KOOS did now not variety between the 2 businesses.

**Table 2 Postoperative variety of motion, knee stability, and final results rating in repair and staged groups**  
**From: touchy fundamental restore of extraarticular ligaments**

**and thespian surgical treatment in multiple ligament knee injuries**

	Repair group (N = 12)	Staged group (N = 19)	p-Value
<b>ROM</b>			
Extension (°): mean ± SD (range)	-2.9 ± 3.2 (-10-0)	-1.4 ± 3.3 (-10-0)	0.104
Flexion (°): mean ± SD (range)	132.5 ± 16.3 (110-150)	134.4 ± 11.7 (105-150)	0.950
<b>Knee stability</b>			
Positive Lachman (≤ grade 2): N (%)	3 (25.0%)	1 (5.3%)	0.279
Positive posterior drawer (≤ grade 2)	9 (75.0%)	2 (10.5%)	0.006
Varus instability (grade 2): N (%)	1(8.3%)	1 (5.3%)	1.000
Valgus instability (grade 2): N (%)	1 (8.3%)	0 (0%)	0.387
<b>Clinical score</b>			
Preinjury Tegner score: mean (range)	4.1 (1-7)	4.7 (1-8)	0.346
Postop Tegner scale: mean (range)	3.3 (1-6)	4.0 (1-8)	0.491
Lysholm score: mean ± SD (range)	87.4 ± 18.5 (44-100)	84.9 ± 19.1 (39-100)	0.537
<b>KOOS: mean ± SD (range)</b>			
Pain	77.3 ± 21.0 (36.1-100)	74.8 ± 21.5 (36.1-100)	0.827
Symptom	75.6 ± 20.6 (42.9-100)	70.2 ± 17.2 (39.3-92.9)	0.610
ADL	79.0 ± 16.3 (57.4-100)	83.5 ± 23.0 (35.3-100)	0.294
Sport/rec	54.5 ± 36.6 (0-100)	57.0 ± 34.8 (10-100)	0.680
QOL	59.7 ± 28.3 (25-100)	61.3 ± 30.1 (25-100)	0.680

1SD trendy deviation, N number, KOOS Knee Injury and Osteoarthritis and Outcome Score, ROM range of motion, ADL sports of day by day living, QOL pleasant of existence

**Discussion:**

surgical field and allows the repair of broken tissue. The maximum important locating of this have a look at is that acute primary restore of the extraarticular ligament provided awesome outcomes for every KDs and MLKIs, and practically comparable consequences have been received for the staged surgical treatment with none detrimental effects. In this collection, 12 of 37 sufferers (38.7%) did no longer need 2d cruciate ligament reconstruction and were glad with their results. Bin and Nam assessed the consequences of the two-diploma manage of their MLKI sufferers and similarly stated that one-1/three of sufferers did now not require 2nd-diploma surgery [18]. They concluded that 2d-level surgical procedure emerge as handiest achieved in instances where it become deemed crucial. Based on those effects, extraarticular ligament repair might also make certain minimal crucial knee balance. Several authors have pronounced systematic reviews of staged surgical treatment for MLKIs [6, 24]. Based on surgical timing, Mook et al. [24] and Jianetal. [6] labeled MLKI remedies into 3 corporations: acute (ligamentous surgical treatment finished much much less than 3 weeks after harm), chronic (ligamentous surgical operation carried out extra than three weeks after damage), and staged (each acute and chronic surgery). They concluded that staged surgical remedy yielded the first-rate clinical effects for MLKIs, no matter the truth that no awesome difference become observed between the intense and persistent surgery groups in medical outcomes. Similarly, Mook et al. Tested that patients who have been controlled acutely had extra flexion deficits than individuals who have been managed chronically. They counseled that extra aggressive rehabilitation may also save you ROM deficits from taking place in acutely handled MLKIs. In other systematic critiques evaluating early as opposed to past due surgical treatment of MLKIs [5, 25], early surgical treatment showed a significantly advanced clinical very last results in contrast with past

due reconstruction. Hohmann et al. Said that general ROM did not significantly differ the various two companies. The MCL and PMC, together with the posterior indirect ligament, are the maximum commonly injured structures in MLKIs [26]. The PMC controls valgus and inner rotation as well as posterior drawer in extension [27]. Therefore, the PMC have to be handled as it must be with the broken MCL. The MCL and PMC can be treated with both number one restore or reconstruction [7]. Since the awesome of the damaged medial systems is commonly strong enough to facilitate a exceptional restore [26], the ones structures need to be repaired all through the extreme section. A systematic assessment of medial knee ligament injuries hooked up that repair of the MCL and PMC become an effective and reliable remedy [28]. Primary repair of these structures advanced now not exceptional valgus balance but moreover affected individual-noted sensible rankings with low fees of secondary failure. An acute primary repair can also preserve grafts for later staged surgical treatment of the cruciate ligament. The PLC is crucial to manipulate varus and rotational stability of the knee, and PLC injuries have a higher incidence than formerly pronounced [29]. Since Stannard et al. Demonstrated that consequences with repair found by using way of early motion rehabilitation were significantly inferior as compared with consequences from reconstruction [30], PLC reconstruction has emerge as a greater well-known technique than number one repair. Similarly, Levy et al. Encouraged reconstruction of the PLC structures primarily based on their comparative cohort take a look at [31]. PLC harm once in a while consists of femoral peel-off lesions that can be efficaciously controlled with primary restore [32]. In this series, repair of the PLC shape showed exceptional results, seemingly because of the truth they had been all repaired in the extreme section, and current suture anchors can be used. Supposedly, the critical issue issue for successful treatment of MLKIs is to keep a proper positional dating among the femur and tibia in knee extension shortly after harm. Since posterior systems, which incorporates the PMC and PLC, come to be taut with knee extension, these structures have a critical function in stabilizing the knee in an extension position. Furthermore, the posterior tablet and indirect popliteal ligament (OPL) are pretty robust structures that contribute to the stability of knee extension [33]. Therefore, those structures want to be in addition repaired. However, present day-day surgical strategies generally overlook about the restore of the posterior pill or the OPL no matter their being willing structures in MLKIs. One study referred to that popliteal artery damage related to MLKI considerably decreased knee characteristic scores compared with those without vascular involvement [34]. However, most research on MLKIs exclude patients with (on three or greater ligaments) had been associated with postoperative stiffness. In this collection, most patients have to accumulate whole knee extension. Since posterior structures of the knee grow to be tight in knee extension, the ones systems need to now not be steady in the flexion feature. Supposedly, this is the reason why no extension deficit become found in our patients. It is essential to check whether or not the knee is truly prolonged after number one restore. In the remedy of acute KDs and MLKIs, accurate prognosis and most appropriate number one repair of extraarticular ligaments are critical to a success control. Extraarticular ligaments must be repaired wherein possible in the extreme phase within the remedy of KDs and MLKIs. This treatment strategy can reduce the frequency of next reconstructive surgery. Evidently, it's miles critical to have a take a look at lengthy-term results, together with development of osteoarthritis, after the ones treatments.

**Limitations:**

We renowned a few barriers to this look at. The maximum enormous trouble is the nonrandomized look at layout. To compare staged surgical remedy and one-stage reconstruction, it might be critical to perform a randomized control trial (RCT). Since MLKIs embody a small cohort with heterogeneous affected person populations, accurate RCTs, as in ACL reconstruction, might be pretty hard to conduct. That number one restore of extraarticular structures improves very last results after cruciate ligament reconstruction in KD or MLKI can only be defined in a comparative look at with a control organization of patients who did no longer go through primary repair. The same is authentic for the proposed discount requiring cruciate ligament reconstruction. Furthermore, it's miles difficult to carry out multiligament reconstruction, which includes the extraarticular ligaments, because of the reality we can't reap allografts in our U.S.A.. The sample length modified into not massive sufficient to reveal the effectiveness of the primary restore. However, the scientific

consequences were similar to the ones mentioned with the useful resource of Bin and Nam [18]. To accumulate 80% statistical electricity with an  $\alpha$  of 0.05 in demonstrating a huge impact period ( $r = 0.5$ ), electricity evaluation discovered that now not less than fifty three patients in each organization might be required for detecting any variations in scientific outcomes between the restore and staged businesses the use of the Mann–Whitney U check. Therefore, further multicenter research are needed.

### Conclusions:

We retrospectively in comparison the effects among acute primary restore of extraarticular ligaments and staged surgical remedy in KD and MLKI. All sufferers once more to their sports of daily living and preinjury profession tiers. Approximately 40% of the patients did no longer require similarly surgical procedure and had been actually as glad with their surgical effects because the staged agency. Acute primary repair of extraarticular ligaments gives crucial knee stability with out varus/valgus instability and can lower the need for subsequent cruciate ligament reconstruction.

### References

- Natsuhara KM, Yerasian MG, Cohen JR et al (2014) What is the frequency of vascular injury after knee dislocation? *Clin Orthop Relat Res* 472:2615–2620 Article Google Scholar
- Scheu M, Espinoza GF, Mellado CA, Díaz PA, Garín AF, O'Connell LA (2020) Varus mechanism is associated with high incidence of popliteal artery lesions in multiligament knee injuries. *Int Orthop* 44:1195–1200 Article Google Scholar
- Darcy G, Edwards E, Hau R (2018) Epidemiology and outcomes of traumatic knee dislocations: isolated vs multi-trauma injuries. *Injury* 49:1183–1187 Article Google Scholar
- Chowdhry M, Burchette D, Whelan D, Nathens A, Marks P, Wasserstein D (2020) Knee dislocation and associated injuries: an analysis of the American College of Surgeons National Trauma Data Bank. *Knee Surg Sports Traumatol Arthrosc* 28:568–575 Article Google Scholar
- Levy BA, Dajani KA, Whelan DB et al (2009) Decision making in the multiligament-injured knee: an evidence-based systematic review. *Arthroscopy* 25:430–438 Article Google Scholar
- Jiang W, Yao J, He Y, Sun W (2015) The timing of surgical treatment of knee dislocations: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 23:3108–3113 Article Google Scholar
- Ng JWG, Myint Y, Ali FM (2020) Management of multiligament knee injuries. *EFORT Open Rev* 5:145–155 Article Google Scholar
- Liow RYL, McNicholas MJ, Keating JF, Nutton RW (2003) Ligament repair and reconstruction in traumatic dislocation of the knee. *J Bone Joint Surg Br* 85:845–851 CAS Article Google Scholar
- Gauffin H, Rockborn P (2014) Knee dislocations: is reconstruction of the posterior cruciate ligament crucial? *Eur J Orthop Surg Traumatol* 24:371–377 Article Google Scholar
- Ibrahim SA, Ahmad FH, Salah M et al (2008) Surgical management of traumatic knee dislocation. *Arthroscopy* 24:178–187 Article Google Scholar
- Li X, Liu T (2013) Surgical management of multiple knee ligament injuries. *Eur J Orthop Surg Traumatol* 23:691–697 Article Google Scholar
- Gwathmey FW, Shafigue DA, Miller MD (2010) Our approach to the management of the multiple-ligament knee injury. *Oper Tech Sports Med* 18:235–244 Article Google Scholar
- LaPrade RF, Chahla J, DePhillipo NN et al (2019) Single-stage multiple-ligament knee reconstructions for sports-related injuries: outcomes in 194 patients. *Am J Sports Med* 47:2563–2571 Article Google Scholar
- Hanley J, Westermann R, Cook S et al (2017) Factors associated with knee stiffness following surgical management of multiligament knee injuries. *J Knee Surg* 30:549–554 Article Google Scholar
- Fanelli GC, Edson CJ (2002) Arthroscopically assisted combined anterior and posterior cruciate ligament reconstruction in the multiple ligament injured knee: 2- to 10-year follow-up. *Arthroscopy* 18:703–714 Article Google Scholar
- Levy BA, Fanelli GC, Whelan DB et al (2009) Knee Dislocation Study Group. Controversies in the treatment of knee dislocations and multiligament reconstruction. *J Am Acad Orthop Surg* 17:197–206 Article Google Scholar
- Moatshe G, Brady AW, Slette EL et al (2017) Multiple ligament reconstruction femoral tunnels: intertunnel relationships and guidelines to avoid convergence. *Am J Sports Med* 45:563–569 Article Google Scholar
- Bin SI, Nam TS (2007) Surgical outcome of 2-stage management of multiple knee ligament injuries after knee dislocation. *Arthroscopy* 23:1066–1072 Article Google Scholar
- van der List JP, DiFelice GS (2017) Primary repair of the anterior cruciate ligament: a paradigm shift. *Surgeon* 15:161–168 Article Google Scholar
- Achtnich A, Herbst E, Forkel P et al (2016) Acute proximal anterior cruciate ligament tears: outcomes after arthroscopic suture anchor repair versus anatomic single bundle reconstruction. *Arthroscopy* 32:2562–2569 Article Google Scholar
- Vermeijden HD, Jonkergouw A, van der List JP, DiFelice GS (2020) The multiple ligament-injured knee: when is primary repair an option? *Knee* 27:173–182 Article Google Scholar
- Schenck RC Jr (1994) The dislocated knee. *Instr Course Lect* 43:127–136 PubMed Article Google Scholar
- Nakamura N, Takeuchi R, Sawaguchi T, Ishikawa H, Saito T, Goldhahn S (2011) Cross-cultural adaptation and validation of the Japanese Knee Injury and Osteoarthritis Outcome Score (KOOS). *J Orthop Sci* 16:516–523 Article Google Scholar
- Mook WR, Miller MD, Diduch DR, Hertel J, Boachie-Adjei Y, Hart JM (2009) Multiple-ligament knee injuries: a systematic review of the timing of operative intervention and postoperative rehabilitation. *J Bone Joint Surg Am* 91:2946–2957 Article Google Scholar
- Hohmann E, Glatt V, Tetsworth K (2017) Early or delayed reconstruction in multiligament knee injuries: a systematic review and meta-analysis. *Knee* 24:909–916 Article Google Scholar
- Tibor LM, Marchant MH Jr, Taylor DC, Hardaker WT Jr, Garrett WE Jr, Sekiya JK (2011) Management of medial-sided knee injuries, part 2: posteromedial corner. *Am J Sports Med* 39:1332–1340 Article Google Scholar
- Petersen W, Loerch S, Schanz S, Raschke M, Zantop T (2008) The role of the posterior oblique ligament in controlling posterior tibial translation in the posterior cruciate ligament-deficient knee. *Am J Sports Med* 36:495–501 Article Google Scholar
- DeLong JM, Waterman BR (2015) Surgical repair of medial collateral ligament and posteromedial corner injuries of the knee: a systematic review. *Arthroscopy* 31:2249–2255 Article Google Scholar
- Feng H, Zhang H, Hong L, Wang XS, Cheng KB, Zhang J (2011) Femoral peel-off lesions in acute posterolateral corner injuries: incidence, classification, and clinical characteristics. *Arthroscopy* 27:951–958 Article Google Scholar
- Stannard JP, Brown SL, Farris RC, McGwin G Jr, Volgas DA (2005) The posterolateral corner of the knee: repair versus reconstruction. *Am J Sports Med* 33:881–888 Article Google Scholar
- Levy BA, Dajani KA, Morgan JA, Shah JP, Dahm DL, Stuart MJ (2010) Repair versus reconstruction of the fibular collateral ligament and posterolateral corner in the multiligament-injured knee. *Am J Sports Med* 38:804–809 Article Google Scholar
- Bonanzinga T, Zhang H, Song GY, Zhang J, Signorelli C, Feng H (2015) Is PLC repair of a peel-off femoral lesion an effective option in a multiligament setting? *Knee Surg Sports Traumatol Arthrosc* 23:2936–2942 Article Google Scholar
- Wu XD, Yu JH, Zou T et al (2017) Anatomical characteristics and biomechanical properties of the oblique popliteal ligament. *Sci Rep* 7:42698 CAS Article Google Scholar
- Sanders TL, Johnson NR, Levy NM et al (2017) Effect of vascular injury on functional outcome in knees with multi-ligament injury: a matched-cohort analysis. *J Bone Joint Surg Am* 99:1565–1571 Article Google Scholar